



**Filed as *Cyperus cuspidatus* Kunth [family
CYPERACEAE]**

<http://plants.jstor.org/stable/10.5555/al.ap.specimen.hnbu00863>

JSTOR

<http://plants.jstor.org>

Your use of JSTOR indicates your acceptance of the Terms & Conditions of Use, available at [http://
www.jstor.org/page/info/about/policies/terms.jsp](http://www.jstor.org/page/info/about/policies/terms.jsp)

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Filed as *Cyperus cuspidatus* Kunth [family CYPERACEAE]

Herbarium	Herbier National du Burkina, Institut de l'Environnement et de Recherche Agricola (HNBU), HNBU00863
Collection	Herbarium Specimens
Resource Type	Specimens
Collector	J. E. Madsen, #6144
Collection Date	1996/10/23
Locality	Tenkodogo. Some kilometers SE of Tenkodogo
Country	Burkina Faso (Burkina Faso)
Collection altitude	300 meters
Identifications	Cyperus cuspidatus Kunth [family CYPERACEAE] (stored under name); Verified by Laegaard S., 1997
Pages	1
Data last modified	2011-08-09

<http://plants.jstor.org/stable/10.5555/al.ap.specimen.hnbu00863>

GP1
2009-2013



FLORA OF BURKINA FASO

J. E. Madsen

6144

Cyperus cuspidatus H. B. K.

Cyperaceae

det. Lægård, S. (AAU), 1997.

Province: BOULGOU

Some km SE of Tenkodogo. Young fallow vegetation with *Ptilostigma reticulatum* and *Cassia obtusifolia*. Also small waterholes and streambeds.
(00° 17' W 11° 43' N) Alt. 300 m.

Oct 23, 1996

Caespitose with spreading branches. On dried-up mud.

Laboratoire de Biologie et Ecologie Végétales (OUAGA),
l'Institut de Recherche en Biologie et Ecologie Tropicale (IRBET) &
Department of Systematic Botany, University of Aarhus, Denmark (AAU)

